

Artikel 2

by Pak Haikal Haikal

Submission date: 23-Jul-2023 10:54AM (UTC+0700)

Submission ID: 2135198143

File name: Artikel_Effect_of_curcumin_extract_Haikal.pdf (767.65K)

Word count: 2234

Character count: 12549



Effect of curcuma extract (*Curcuma xanthorrhiza*) on piroxicam-induced damage to the small intestine cells of mice (*Mus musculus*)

Moch. Haikal^{1*}, Septiana Kurniasari²

¹*Biology Education Study Program, Madura Islamic University, Indonesia*

²*D3 Pharmacy Study Program, Madura Islamic University, Indonesia*

*E-mail: moch.haikal@fkip.uim.ac.id

(Received: 04 January 2021; Revised: 14 February 2021; Accepted: 26 February 2021)

ABSTRACT

Piroxicam is a non-steroidal anti-inflammatory drug (NSAID) widely used as an analgesic and anti-inflammatory in rheumatoid arthritis, osteoarthritis. However, Piroxicam can cause gastrointestinal side effects and peptic ulcers when given orally. Piroxicam can also cause cholestasis. Also, if Piroxicam is used continuously in high doses, it can cause damage to the small intestine. A compound can minimize small intestinal cell damage called an antioxidant. This study aims to analyze the effect of curcuma extract (*Curcuma xanthorrhiza*) on Piroxicam-induced damage to small intestinal cells of mice (*Mus musculus*). This study used 80 male mice and divided them into 3 groups, namely negative control (K-), negative Piroxicam (P-), and positive Piroxicam (P+). Giving curcuma extract given orally at a dose of 1.4 g/kgBW; 2.0 g/kgBW; 2.6 g/kgBW; 3.2 g/kgBW and 3.8 g/kgBW, while the Piroxicam dosage used was 10 mg/kgBW and 30 mg/kgBW. The results showed that the more Piroxicam doses were given, the higher the level of small intestinal cell damage. Along with increasing the dosage of curcuma extract, the level of damage to small intestinal cells is reduced.

Keywords: curcuma, piroxicam, small intestine

DOI:[10.30870/gravity.v7i1.10181](https://doi.org/10.30870/gravity.v7i1.10181)

INTRODUCTION

Piroxicam is a non-steroidal anti-inflammatory drug (NSAID) widely used as an analgesic and anti-inflammatory in rheumatoid arthritis, osteoarthritis. However, Piroxicam can cause side effects in the gastrointestinal tract and peptic ulcers when given orally (Aryani, 2007). Piroxicam can also cause cholestasis. Also, if Piroxicam is used continuously in high doses, it can cause damage to small intestinal cells. Antioxidant

compounds can minimize damage to the small intestine cells. One of them is curcuma (*Curcuma xanthorrhiza*). Temulawak contains main components, including curcumin, essential oils, fats, dyes, proteins, resins, cellulose, pentosan, starch, and minerals (Oktaviana, 2010).

Several research results prove that curcuma can potentially be a hepatoprotection in mice injected intraperitoneally with a toxic dose of paracetamol. Also, the content of curcumin functions as an antioxidant and detoxifying

agent from increased activity of the glutathione S-transferase (GS-t) enzyme and other glutathione (GS-x) enzymes and can protect erythrocytes and hemoglobin from oxidation caused by nitrite compounds (Sugiharto, 2004).

Curcuma xanthorrhiza has long been recognized in Asia as an antihepatotoxic. This plant is widely used to treat hepatitis C. Xanthorrhiza plants are effective in reducing hepatotoxicity levels. This is also supported by other studies showing that *C. xanthorrhiza* has a hepatoprotective effect that can act as an effective treatment for acute liver disease in rats (Putri, 2013).

Curcumin can also protect the liver's function, digestive tract, kidneys, reduce free radicals, and inhibit the activity of nitric oxide synthase (NOS) from macrophages (Kertia, 2011). Also, curcuma is also used to increase endurance and stamina (Dewi, et al., 2012). Temulawak can overcome liver disorders, increase bile production and secretion, anti-inflammatory, appetite enhancer, asthma medication, antioxidants, inhibit blood clotting and reduce ALT and SGOT levels (Khamidah, Antarlina & Sudaryono, 2017).

The content in curcuma contains chemical compounds that are physiologically active, for example, curcumin, starch, and essential oils. The content of curcumin in curcuma functions as antibacterial, anti-cancer, anti-tumor, and contains antioxidants. The curcuminoid content in curcuma is 1-2%, and the essential oil content in curcuma is 3-12% (Pratama, Ismail & Witjahjo, 2019).

Temulawak (*Curcuma xanthorrhiza* Roxb.) Is one of the many medicinal plants of the Zingiberaceae tribe that grows in Indonesia. Temulawak is known to have many benefits, including anti-hepatitis, anti-carcinogenic, antimicrobial, antioxidant, anti-hyperlipidemic, antiviral, anti-inflammatory, and detoxifying agent (Candra, 2013). Other properties of curcuma's chemical components are antibacterial, anti-fungal, neuroprotector, anti-cancer, allergy, and anti-hypercholesterolemic (Nurcholis, et al., 2012).

Previous research has proven that curcuma has fungistatic properties against several

dermatophyta fungi and is bacteriostatic in the microbes such as Salmonella, Pseudomonas pyogenes, Staphylococcus aureus, and anti-fungi against *Mycrosporium gypseum*. The activity of 1 ml of essential oil on Pseudomonas pyogenes was comparable to 0.3 mg of tetracyclines. The action of 1 ml of essential oils on Staphylococcus aureus was equal to 29.2658 mg of tetracyclines. It is suspected that the actual oil content in the curcuma rhizome is antibacterial (Mashita, 2014).

The formulas of curcuma, turmeric, and meniran jamu can improve the tested animals' fitness as indicated by the increase in swimming endurance time of the test animals after treatment compared to the pre-treatment and control groups. The herbal formula of curcuma, turmeric, and meniran has also been proven to be safe for the liver and kidney function of test animals given doses of up to 2,160 mg/kg bodyweight for 90 consecutive days (Novianto, et al., 2020).

RESEARCH METHODS

This study used 80 male mice with an average body weight of 18-20 grams. The mice were grouped into 3 groups, namely K- (without Piroxicam and extract), P- (Piroxicam without extract) and P+ (given Piroxicam then extract). The doses of Piroxicam used were 10 mg/kgBW and 30 mg/kgBW, while temulawak extract was given in five dose variations, namely 1.4 g/kgBW; 2.0 g/kgBW; 2.6 g/kgBW; 3.2 g/kgBW and 3.8 g/kgBW.

Giving of piroxicam. Piroxicam given to mice is a finished powder that has been packaged and sold in the market, taking into account the composition contained therein. One Piroxicam capsule contains 10 mg. The dose of Piroxicam given to mice is calculated based on the bodyweight of each mouse. Piroxicam was given once a day for 7 days before the mice were given curcuma extract by force-feeding the mice using a gastric swab.

Giving of Temulawak Extract. Temulawak extract given to mice is a finished powder that has been packaged and sold in the market, taking into account the composition contained

therein. One curcuma extract capsule contains 500 mg of curcumin. The dose of curcuma extract given to mice was calculated based on the bodyweight of each mouse. Temulawak extract was shown once a day for 7 days after the mice were given Piroxicam by feeding it to the mice using a gastric swab.

Preparation of Histology. For mice that have been operated on, their small intestine organs are taken. The small intestine of mice was dehydrated and then cut with a microtome. The pieces of the small intestine were rehydrated, and HE stained. Furthermore, it can be observed under a microscope.

RESULTS AND DISCUSSION

The data obtained in this study were the width of the villi that had broken off and indicated damage to the small intestine cells of mice under normal conditions, was induced by Piroxicam with 2 variations of doses, and induced by Piroxicam then given curcuma extract.




Figure 1. The Relationship of the Number of Piroxicam Doses to the Dilatation of Villi in Mice Small Intestine Cells

Figure 1 shows a graph of the relationship between the number of Piroxicam doses on the enlargement of the villi in mice's small intestine cells. The amount of Piroxicam in mice with 2 dosage variations, namely 10 mg/kgBW and 30 mg/kgBW, indicates that the more Piroxicam doses are given, the greater the enlargement of the villi. This suggests that the damage to the small intestine cells of mice is getting worse. In the adverse control treatment, the width of the villi was 6.21 µm. In the Piroxicam treatment of 10 mg/kgBW and 30 mg/kgBW, the villi's widths were 10.20 µm and 15.20 µm, respectively.




Figure 2. Microscopic picture of small intestinal cells of mice (a) control (b) Piroxicam 10 mg/kgBW (c) Piroxicam 30 mg/kgBW

Figure 2 shows a microscopic view of small intestinal cells of mice taken at 400x magnification. In control mice (standard), shown in Figure 2a, the severed villi were not very wide. However, the more doses of Piroxicam given, the greater the dilation or rupture of the villi that occurs. Giving curcuma extract also greatly affects the widening of the villi in the small intestine of mice. This is shown in Figure 3, and where there was a decrease in villi width in mice, both treated with Piroxicam dose 10 mg/kg BW and Piroxicam dose 30 mg/kg BW.





Figure 3. The Relationship of the Number of Doses of Temulawak Extract on the Dilatation of Villi in Mice Small Intestine Cells



REFERENCES

- Aryani, N. L. D. S. M. (2007). Uji Permeabilitas Intrinsik dan Termodinamika Difusi Piroksikam secara *In Vitro*. *Farmasi Indonesia*, 3, 103-110.
- Candra, A. A. (2013). Aktivitas Hepatoprotektor Temulawak pada Ayam yang Diinduksi Pemberian Parasetamol. *Jurnal Penelitian Pertanian Terapan*, 13 (2), 137-143.
- Dewi, dkk. (2012). Pengetahuan tentang Manfaat Kesehatan Temulawak (*Curcuma xanthorrhiza*) serta Uji Klinis Pengaruhnya pada Sistem Imun Humoral pada Dewasa Obes. *Jurnal Ilmu Pertanian Indonesia*, 17(3), 166-171.
- Kertia, N. (2011). Pengaruh Kombinasi Ekstrak Temulawak, Jahe, Kedelai dan Kulit Udang terhadap Fungsi Hati dan Ginjal Dibandingkan dengan Natrium Diklofenak pada Penderita Osteoarthritis. *Majalah Obat Tradisional*, 16, 51-62.
- Khamidah, A., Antarlina, S. S., & Sudaryono, T. (2017). Ragam Produk Olahan Temulawak untuk Mendukung Keanekaragaman Pangan. *Jurnal Litbang Pertanian*, 36(1), 1-12.
- Mashita, A. R. (2014). Efek Antimikroba Ekstrak Rimpang Temulawak (*Curcuma xanthorrhiza*) terhadap Pertumbuhan. 10 (2), 138-144.
- Novianto, dkk. (2020). Pengaruh Formula Jamu Temulawak, Kunyit, dan Meniran terhadap Kebugaran Jasmani: Suatu Studi Klinik. *Media Litbangkes*, 30(1), 37-44.
- Nurcholis, dkk. (2012). Variasi Bahan Bioaktif dan Bioaktivitas Tiga Nomor Harapan Temulawak pada Lokasi Budidaya Berbeda. *J. Agron. Indonesia*, 40(2), 153-159.
- Oktaviana, P. R. (2010). Kajian Kadar Kurkuminoid, Total Fenol dan Aktivitas Antioksidan Ekstrak Temulawak (*Curcuma xanthorrhiza* Roxb.) pada Berbagai Teknik Pengeringan dan Proporsi Pelarutan. Surakarta: 86.
- Pairul, P. P. B. (2018). Perbedaan Efek Anti Inflamasi Jahe Merah (*Zingiber officinale* Rosc. Var. *Rubrum*) dan Jahe Putih Besar (*Zingiber officinale* Rosc. Var. *Officinarum*) terhadap Ulkus Gaster Tikus Jantan Galur *Sprague dawley* yang Diinduksi Piroksikam. Skripsi. Bandar Lampung, Universitas Lampung. S1: 65.
- Pratama, P. B., Ismail, A., & Witjahjo, R. B. B. (2019). Pengaruh Pemberian Ekstrak Temulawak (*Curcuma xanthorrhiza*) Dosis Bertingkat terhadap Gambaran Mikroskopis Hepar Mencit Balb/C Jantan yang Diinduksi Rifampisin. *JURNAL KEDOKTERAN DIPONEGORO*, 8(3), 1026-1036.
- Putri, R. M. S. (2013). Si “Kuning” Temulawak (*Curcuma xanthoriza* Roxb.) dengan “Segudang” Khasiat. *Jurnal Teknologi Pertanian*, 2(2), 42-49.
- Ramadhan, R. I. (2015). Rasionalitas Penggunaan OAINS pada Pasien Rematik Osteoarthritis Rawat Jalan di RSUD Kabupaten Subang Tahun 2014 Ditinjau dari Tepat Diagnosis, Tepat Indikasi, Tepat Obat, Tepat Dosis, Tepat Cara Pemberian, Tepat Pasien. Skripsi. Jakarta, UIN Syarif Hidayatullah Jakarta. S1: 141.
- Sari, S. K., Widodo, C. S., & Juswono, U. P. (2018). Efek Paparan Radiasi Gamma dan Pemberian Ekstrak Temulawak (*Curcuma xanthorrhiza*) terhadap Pelebaran Vena Centralis Hepar Mencit (*Mus musculus*). *Journal of Medical Physics and Biophysics*, 5(2), 203-210.
- Sugiharto. (2004). Pengaruh Infus Rimpang Temulawak (*Curcuma xanthorrhiza*) terhadap Kadar Hemoglobin dan Jumlah Eritrosit Tikus Putih yang Diberi Larutan Timbal Nitrat [(PbNO₃)₂]. *Hayati*, 10, 53-57.

Artikel 2

ORIGINALITY REPORT

11 %

SIMILARITY INDEX

11 %

INTERNET SOURCES

3 %

PUBLICATIONS

5 %

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

4%

★ jurnal.untirta.ac.id

Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography On