

The local strong metric dimension in the join of graphs

by Rica Dkk

Submission date: 16-Dec-2022 09:22AM (UTC+0700)

Submission ID: 1982484050

File name: 9_Amalia_2022_J_Phys._Conf._Ser._2157_012004.pdf (1.09M)

Word count: 2586

Character count: 11839

PAPER • OPEN ACCESS

The local strong metric dimension in the join of graphs

To cite this article: R Amalia *et al* 2022 *J. Phys.: Conf. Ser.* **2157** 012004

View the [article online](#) for updates and enhancements.

You may also like

- [On the metric dimension of generalized broken fan graph and edge corona product of path with star graph](#)
Makarima Sabila and Tri Almojo Kusmayadi
- [Metric and edge-metric dimensions of bobble-neighbourhood-corona graphs](#)
Rinuwati and R E Nabila
- [Fault-Tolerant Metric Dimension of Cube of Paths](#)
Laxman Saha



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII
Boston, MA • May 28 – June 2, 2023

**Abstract Submission Extended
Deadline: December 16**

[Learn more and submit!](#)

The local strong metric dimension in the join of graphs

R Amalia¹, Firdausiyah¹, T Yulianto¹, Faisal¹ and Kuzairi¹

¹Mathematics Department, Mathematics and Science Faculty, Universitas Islam Madura Pamekasan, Indonesia

E-mail: rica.amalia@uim.ac.id

Abstract. Let G be a connected graph. A vertex w is said to strongly resolve a pair u, v of vertices of G if there exists some shortest $u - w$ path containing v or some shortest $v - w$ path containing u . A set W of vertices is a local strong resolving set for G if every pair of adjacent vertices of G is strongly resolved by some vertex of W . The smallest cardinality of a local strong resolving set for G is called the local strong metric dimension of G . In this paper we studied the local strong metric dimension in the join of graphs. We use the path, cycle, complete, and star graphs in this studies.

1. Introduction

Graph theory is a subject in mathematics which first introduced by a Swiss mathematician named Leonard Euler in 1736, as an effort to solve the Konigsberg Bridge problem [4]. Graphs are used to represent discrete objects and the relationships between those objects. The visual representation of a graph is to represent objects as vertices (nodes) and the relationships between objects as edges [2]. One of the studies that continue to develop in graph theory is metric dimension.

The concept of metric dimension in graph theory has been introduced separately by Slater in 1975 and by Harrary and Melter in 1976. Slater relates the problem of metric dimensions to determine the number of sonar detection tools in a network [8] while Harrary defines metric dimensions through the set of differentiators. This concept can be used to distinguish each point on a graph G by determining its representation against the set of vertices from G [1]. The concept of metric dimension then was developed into local metric dimensions by Okamoto et al. [7]. This concept states that every adjacent vertices on a graph has different representation in regard to local differentiators [3]. Okamoto et al. obtained the result that a nontrivial connected graph G has local metric dimension $n - 1$ if and only if G is complete graphs. Also, graph G has a local metric dimension of 1 if and only if G is bipartite graphs.

Another developed concept of metric dimension is strong metric dimension. This concept is found by Sebo and Tannier in 2004. A vertex w is said to strongly resolve a pair u, v of vertices of G if there exists some shortest $u - w$ path containing v or some shortest $v - w$ path containing u [6]. A set that contains those vertices is called strong resolving set of G . The strong metric dimension of G is the smallest cardinality of strong resolving set, denoted by $sdim(G)$. The concept of strong metric dimension then developed into local strong metric dimension [10]. A set W of vertices is a local strong resolving set for G if every pair of adjacent vertices of G is strongly resolved by some vertex of W . The smallest cardinality of a local strong resolving set for G is called the local strong metric dimension of G , denoted by $dim_{ls}(G)$. Several results regarding local strong metric dimension has been found in path graph, star graph, complete graph, cycle graph, and graph resulting from corona product.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

Research on local strong metric dimension has been carried out by Susilowati et al. [9] on k -level corona product graphs. The next research was conducted by Sutardji et al. [10] on cartesian product graph. Furthermore, research on the graph resulting from join operation has been carried out by Kuziak et al. [5] to get the strong metric dimension.

Therefore, in this paper we discuss the local strong metric dimension in the join of graphs.

Theorem 1.1 [10] *Let G be a connected graph of order $n \geq 2$, then*

- i. $\dim_{ls}(G) = 1$ if and only if G is bipartite graph
- ii. $\dim_{ls}(G) = n - 1$ if and only if G is complete graph

Theorem 1.2 [9] *Let P_n be a path graph, C_n be a cycle, and S_n be a star graph, then*

1. $\dim_{ls}(P_n) = 1$
2. $\dim_{ls}(C_n) = \begin{cases} 1 & \text{if } n \text{ is even} \\ 2 & \text{if } n \text{ is odd} \end{cases}$
3. $\dim_{ls}(S_n) = 1$

2. Local strong metric dimension of $K_1 + G$

In this section, we obtained the local strong metric dimension in the join between a trivial graph (K_1) and G , where G is one of the following graphs, which are path (P_n), cycle (C_n), complete graph (K_n), and star graph (S_n).

Before we discuss the local strong metric dimension in those graphs, we need to know the symbol of $I[u, v]$ which denotes the shortest path between vertex u and vertex v . If $W \subseteq V(G)$ is a local strong resolving set, then a vertex $w \in W$ is said to strongly resolve an adjacent pair u, v of vertices of G if $u \in I[v, w]$ or $v \in I[u, w]$. We do not have to check every adjacent vertices in G , just adjacent vertices in G other than W .

Generally speaking, if G is graph resulting from join operation then $diam(G) = 2$, where $diam(G)$ is diameter of G which is $diam(G) = \max_{u, v \in V(G)} d(u, v)$. So, in the join of graphs, $d(u, v) = 2$ if u and v are not adjacent vertices.

Theorem 2.1 *Let G be a join graph $K_1 + P_n$, where P_n is a path, then $\dim_{ls}(G) = 2$ if $2 \leq n \leq 5$ and*

$$\dim_{ls}(G) = \begin{cases} \left\lfloor \frac{n-3}{4} \right\rfloor + 1 & \text{for } (n-3) \bmod 4 < 3 \\ \left\lfloor \frac{n-3}{4} \right\rfloor + 1 & \text{for } (n-3) \bmod 4 = 3 \end{cases} \text{ if } n > 5, \text{ where } [x] = m \text{ for } m < x < m + 1 \text{ and}$$

m is integer.

Proof.

Let $V(G) = \{c\} \cup \{v_i | i = 1, 2, \dots, n\}$ and $E(G) = \{cv_i | 1 \leq i \leq n\} \cup \{v_i v_{i+1} | i = 1, 2, \dots, n-1\}$.

Case 1. For $2 \leq n \leq 5$

Let $W = \{c, v_{\frac{n+1}{2}}\}$ if n is odd and $W = \{c, v_{\frac{n}{2}}\}$ if n is even. Then for every pair of adjacent vertices in $V(G) \setminus W$:

- i. $(v_i, v_{i+1}) \rightarrow v_{i+1} \in I[v_i, v_{i+2}]$ or $v_i \in I[v_{i-1}, v_{i+1}]$
- ii. $(c, v_i) \rightarrow c \in I[v_i, v_j]$ where v_i and v_j are not adjacent

So, W is a local strong resolving set of G .

Next, we will prove that W is the local strong resolving set with the smallest cardinality. Take any set $W' \subseteq V(G)$ where $|W'| < 2$, so there are two option of W' which are:

- i. If $W' = \{c\}$ the 1 for a pair adjacent vertice $(v_i, v_{i+1}) \rightarrow v_{i+1} \notin I[v_i, c]$ and $v_i \notin I[v_{i+1}, c]$. So $W' = \{c\}$ is not a local strong resolving set of G .
- ii. If $W' = \{v_i\}$ the 1 for a pair adjacent vertice $(c, v_{i+1}) \rightarrow v_{i+1} \notin I[c, v_i]$ and $c \notin I[v_{i+1}, v_i]$. So $W' = \{v_i\}$ is not a local strong resolving set of G .

Based on the argument above, W' where $|W'| = 1$ is not a local strong resolving set of G . Therefore, W is the local strong resolving set with the smallest cardinality. Thus, $\dim_{ls}(G) = 2$ if $2 \leq n \leq 5$.

Case 2. For $n > 5$

Let $W = \{v_{4i+3} | i = 0, 1, \dots, \lfloor \frac{n-3}{4} \rfloor\}$ if $(n-3) \bmod 4 < 3$ and $W = \{v_{4i+3} | i = 0, 1, \dots, \lfloor \frac{n-3}{4} \rfloor\} \cup \{v_n\}$ if $(n-3) \bmod 4 < 3$. Then for every pair of adjacent vertices in $V(G) \setminus W$:

- i. $(v_1, v_2) \rightarrow v_2 \in I[v_1, v_3]$
- ii. $(v_{4i+4}, v_{4i+5}) \rightarrow v_{4i+4} \in I[v_{4i+5}, v_{4i+3}]$ for $i = 0, 1, \dots, \lfloor \frac{n-3}{4} \rfloor$
- iii. $(v_{4i+5}, v_{4i+6}) \rightarrow v_{4i+6} \in I[v_{4i+5}, v_{4(i+1)+3}]$ for $i = 0, 1, \dots, \lfloor \frac{n-3}{4} \rfloor$

So, W is a local strong resolving set of G .

Next, we will prove that W is the local strong resolving set with the smallest cardinality. Take any set $W' \subseteq V(G)$ where $|W'| < |W|$. Suppose $W' = W - \{a\}$ where $a \in W$, then:

- i. If $W' = W - \{v_3\}$ then for a pair adjacent vertice $(v_1, v_2) \rightarrow v_2 \notin I[v_1, w]$ and $v_1 \notin I[v_2, w]$, where $w \in W'$. So W' is not a local strong resolving set of G .
- ii. If $W' = W - \{v_{4k+3}\}$ where $k = 1, 2, \dots, \lfloor \frac{n-3}{4} \rfloor$ then for a pair adjacent vertice $(v_{4k+1}, v_{4k+2}) \rightarrow v_{4k+1} \notin I[v_{4k+2}, w]$ and $v_{4k+2} \notin I[v_{4k+1}, w]$, where $w \in W'$. So W' is not a local strong resolving set of G .

Based on the argument above, W' where $|W'| < |W|$ is not a local strong resolving set of G . Therefore, W is the local strong resolving set with the smallest cardinality. Thus, $\dim_{ls}(G) =$

$$\begin{cases} \lfloor \frac{n-3}{4} \rfloor + 1 & \text{for } (n-3) \bmod 4 < 3 \\ \lfloor \frac{n-3}{4} \rfloor + 1 & \text{for } (n-3) \bmod 4 = 3 \end{cases} \text{ if } n > 5. \quad \blacksquare$$

Theorem 2.2 Let G be a join graph $K_1 + C_n$, where C_n is a cycle graph, then $\dim_{ls}(G) = 3$ if $3 \leq n \leq 5$ and $\dim_{ls}(G) = \lfloor \frac{n-1}{4} \rfloor + 1$ if $n > 5$, where $\lfloor x \rfloor = m$ for $m < x < m + 1$ and m is integer.

Proof.

Let $V(G) = \{c\} \cup \{v_i | i = 1, 2, \dots, n\}$ and $E(G) = \{cv_i | i = 1, 2, \dots, n\} \cup \{v_i v_{i+1} | i = 1, 2, \dots, n-1\} \cup \{v_1 v_n\}$.

Case 1. For $3 \leq n \leq 5$

Let $W = \{c, v_1, v_{\frac{n+1}{2}}\}$ if n is odd and $W = \{c, v_1, v_{\frac{n}{2}}\}$ if n is even. Then for every pair of adjacent vertices in $V(G) \setminus W$:

- i. $(v_1, v_n) \rightarrow v_1 \in I[v_n, v_2]$ or $v_n \in I[v_1, v_{n-1}]$
- ii. $(v_i, v_{i+1}) \rightarrow v_{i+1} \in I[v_i, v_{i+2}]$ or $v_i \in I[v_{i-1}, v_{i+1}]$
- iii. $(c, v_i) \rightarrow c \in I[v_i, v_j]$ where v_i and v_j are not adjacent

So, W is a local strong resolving set of G .

Next, we will prove that W is the local strong resolving set with the smallest cardinality. Take any set $W' \subseteq V(G)$ where $|W'| < 3$, so there are three option of W' which are:

- i. If $W' = \{c, v_i\}$ then for a pair adjacent vertice $(v_{i+1}, v_{i+2}) \rightarrow v_{i+1} \notin I[v_{i+2}, c]$ and $v_{i+2} \notin I[v_{i+1}, c]$. So $W' = \{c, v_i\}$ is not a local strong resolving set of G .
- ii. If $W' = \{v_i, v_{i+1}\}$ then for a pair adjacent vertice $(c, v_{i+2}) \rightarrow v_{i+2} \notin I[c, v_i]$ and $c \notin I[v_{i+2}, v_{i+1}]$. So $W' = \{v_i, v_{i+1}\}$ is not a local strong resolving set of G .
- iii. If $W' = \{v_i, v_j\}$, where v_i and v_j are not adjacent, then for a pair adjacent vertice $(c, v_{j+1}) \rightarrow v_{j+1} \notin I[c, v_i]$ and $c \notin I[v_{j+1}, v_j]$. So $W' = \{v_i, v_j\}$ is not a local strong resolving set of G .

Based on the argument above, W' where $|W'| = 2$ is not a local strong resolving set of G . Therefore, W is the local strong resolving set with the smallest cardinality. Thus, $\dim_{ls}(G) = 3$ if $3 \leq n \leq 5$.

Case 2. For $n > 5$

Let $W = \{v_{4i+1} | i = 0, 1, \dots, \lfloor \frac{n-1}{4} \rfloor\}$. Then for every pair of adjacent vertices in $V(G) \setminus W$:

- i. $(v_{4i+2}, v_{4i+3}) \rightarrow v_{4i+2} \in I[v_{4i+3}, v_{4i+1}]$ for $i = 0, 1, \dots, \lfloor \frac{n-1}{4} \rfloor$
- ii. $(v_{4i+3}, v_{4i+4}) \rightarrow v_{4i+4} \in I[v_{4i+3}, v_{4(i+1)+1}]$ for $i = 0, 1, \dots, \lfloor \frac{n-1}{4} \rfloor$

iii. $(c, v_i) \rightarrow c \in I[v_i, v_{4i+1}]$ for $i = 1, 2, \dots, \lfloor \frac{n-1}{4} \rfloor$

So, W is a local strong resolving set of G .

Next, we will prove that W is the local strong resolving set with the smallest cardinality. Take any set $W' \subseteq V(G)$ where $|W'| < |W|$. Suppose $W' = W - \{a\}$ where $a \in W$, then if $W' = W - \{v_{4k+1}\}$ where $k = 0, 1, \dots, \lfloor \frac{n-1}{4} \rfloor$ then for a pair adjacent vertex $(v_{4k+2}, v_{4k+3}) \rightarrow v_{4k+2} \notin I[v_{4k+1}, w]$ and $v_{4k+3} \notin I[v_{4k+2}, w]$, where $w \in W'$. So W' is not a local strong resolving set of G .

Based on the argument above, W' where $|W'| < |W|$ is not a local strong resolving set of G . Therefore, W is the local strong resolving set with the smallest cardinality. Thus, $\dim_{ls}(G) = \lfloor \frac{n-1}{4} \rfloor + 1$ if $n > 5$. ■

Theorem 2.3 Let G be a join graph $K_1 + K_n$, where K_n is a complete graph with order $n \geq 2$, then $\dim_{ls}(G) = n$

Proof.

Let $V(G) = \{c\} \cup \{v_i | 1 \leq i \leq n\}$ and $E(G) = \{cv_i | i = 1, 2, \dots, n\} \cup \{v_i v_j | i, j = 1, 2, \dots, n\}$.

Graph $G = K_1 + K_n$ is isomorphic with graph K_{n+1} . Based on Theorem 1.1, $\dim_{ls}(K_{n+1}) = n + 1 - 1 = n$. Thus $\dim_{ls}(G) = n$. ■

Theorem 2.4 Let G be a join graph $K_1 + S_n$, where S_n is a star graph with order $n \geq 2$, then $\dim_{ls}(G) = 2$

Proof.

Let $V(G) = \{c\} \cup \{v_i | i = 0, 1, \dots, n\}$ and $E(G) = \{cv_i | i = 0, 1, \dots, n\} \cup \{v_0 v_i | i = 1, 2, \dots, n\}$.

Suppose $W = \{c, v_0\}$, then for every pair of adjacent vertices in $V(G)$:

i. $(c, v_i) \rightarrow c \in I[v_i, c]$ for $i = 0, 1, \dots, n$

ii. $(v_0, v_i) \rightarrow v_0 \in I[v_i, v_0]$ for $i = 1, 2, \dots, n$

So, W is a local strong resolving set of G .

Next, we will prove that W is the local strong resolving set with the smallest cardinality. Take any set $W' \subseteq V(G)$ where $|W'| < 2$, so there are three option of W' which are:

i. If $W' = \{c\}$ then for a pair adjacent vertex $(v_0, v_i) \rightarrow v_i \notin I[v_0, c]$ and $v_0 \notin I[v_i, c]$. So $W' = \{c\}$ is not a local strong resolving set of G .

ii. If $W' = \{v_0\}$ then for a pair adjacent vertex $(c, v_i) \rightarrow v_i \notin I[c, v_0]$ and $c \notin I[v_i, v_0]$. So $W' = \{v_0\}$ is not a local strong resolving set of G .

iii. If $W' = \{v_k\}$ where $k = 1, 2, \dots, n$ then for a pair adjacent vertex $(c, v_0) \rightarrow v_0 \notin I[c, v_k]$ and $c \notin I[v_0, v_k]$. So $W' = \{v_k\}$ is not a local strong resolving set of G .

Based on the argument above, W' where $|W'| = 1$ is not a local strong resolving set of G . Therefore, W is the local strong resolving set with the smallest cardinality. Thus, $\dim_{ls}(G) = 2$. ■

3. Local strong metric dimension of $G + H$

In this section, we get the local strong metric dimension in the join of graphs, $G + H$, with G and H be a connected graph. First, we discuss the properties of the local strong basis of $G + H$. Local strong basis is the local strong resolving set with the minimum cardinality and this cardinality is called the local strong metric dimension.

Lemma 3.1 Let $W = W_1 \cup W_2$ be the local strong basis of $G + H$ where $W_1 \subseteq V(G)$ and $W_2 \subseteq V(H)$, where $|V(G)| \geq 2$ and $|V(H)| \geq 2$, then $\dim_{ls}(G + H) \geq 3$, with $W_1 \neq \emptyset$ and $W_2 \neq \emptyset$.

Proof. Let $V(G) = \{u_i | i = 1, 2, \dots, n\}$ and $V(H) = \{v_i | i = 1, 2, \dots, m\}$. First, we assume that $W_1 = \emptyset$ or $W_2 = \emptyset$. If $W_1 = \emptyset$ then $W = W_2$. So, for a pair of adjacent vertices $(u_i, u_j) \rightarrow u_i \notin I[u_j, v_k]$ and $u_j \notin I[u_i, v_k]$ where $v_k \in W_2$. On the other hand, if $W_2 = \emptyset$ then $W = W_1$. So, for a pair of adjacent vertices $(v_i, v_j) \rightarrow v_i \notin I[v_j, u_k]$ and $v_j \notin I[v_i, u_k]$ where $u_k \in W_1$. Therefore, $W_1 \neq \emptyset$ and $W_2 \neq \emptyset$.

Next, if we let $W = \{u_k, v_k\}$ then for a pair of adjacent vertices (u_i, v_j) where u_i adjacent with u_k and v_i adjacent with v_k applies $v_j \notin I[u_i, u_k]$ and $u_i \notin I[v_j, v_k]$. So, W with two elements is not the local strong basis of $G + H$. Thus $\dim_{ls}(G + H) \geq 3$. ■

Lemma 3.2 Let the join of two connected graphs, $G + H$, with $|V(G)| = n$ and $|V(H)| = m$ and $V(G + H) = \{u_i | i = 1, 2, \dots, n\} \cup \{v_i | i = 1, 2, \dots, m\}$ then $d(u_i, u_j) = d(v_i, v_j) = 2$ if u_i is not adjacent to u_j and v_i is not adjacent to v_j . Furthermore, $d(u_i, v_j) = 1$.

Proof. Since $V(G + H) = \{u_i | i = 1, 2, \dots, n\} \cup \{v_i | i = 1, 2, \dots, m\}$ then $u_i v_j \in E(G + H)$. Thus $d(u_i, v_j) = 1$. Next, since u_i is not adjacent to u_j and v_i is not adjacent to v_j then $d(u_i, u_j) \neq 1$ and $d(v_i, v_j) \neq 1$. Because $diam(G + H) = 2$ then $d(u_i, u_j) = d(v_i, v_j) = 2$. ■

Using the lemma above, we get the local strong metric dimension of $G + H$ which is stated in the following theorem.

Theorem 3.1. Let G be a connected graph with order $n \geq 2$ and H be a connected graph with order $m \geq 2$ then

$$\dim_{ls}(G + H) = \begin{cases} \dim_{ls}(K_1 + G) + \dim_{ls}(K_1 + H) & \text{if } diam(G) > 2 \text{ and } diam(H) > 2 \\ \dim_{ls}(K_1 + G) + \dim_{ls}(H) & \text{if } diam(G) > 2 \text{ and } diam(H) \leq 2 \\ \dim_{ls}(G) + \dim_{ls}(K_1 + H) & \text{if } diam(G) \leq 2 \text{ and } diam(H) > 2 \\ \dim_{ls}(G) + \dim_{ls}(H) + 1 & \text{if } diam(G) \leq 2 \text{ and } diam(H) \leq 2 \\ & \text{or } \dim_{ls}(G) = \dim_{ls}(H) = 1 \end{cases}$$

Proof. Let $V(G) = \{u_i | i = 1, 2, \dots, n\}$, $V(H) = \{v_i | i = 1, 2, \dots, m\}$, $V(G + H) = V(G) \cup V(H)$, and $E(G + H) = E(G) \cup E(H) \cup \{u_i v_j | i = 1, 2, \dots, n; j = 1, 2, \dots, m\}$. Suppose $W = W_1 \cup W_2$ where $W_1 \subseteq V(G)$ and $W_2 \subseteq V(H)$. In order to get the minimum cardinality of W , we get four cases which are:

Case 1. If $diam(G) > 2$ and $diam(H) > 2$

According to Lemma 3.2, $d(u_i, u_j) = d(v_i, v_j) = 2$ if u_i is not adjacent to u_j and v_i is not adjacent to v_j . In order to satisfy this lemma in this case, $|W_1| = \dim_{ls}(K_1 + G)$ and $|W_2| = \dim_{ls}(K_1 + H)$. So, $\dim_{ls}(G + H) = \dim_{ls}(K_1 + G) + \dim_{ls}(K_1 + H)$ if $diam(G) > 2$ and $diam(H) > 2$.

Case 2. If $diam(G) > 2$ and $diam(H) \leq 2$

According to Lemma 3.2, $d(u_i, u_j) = d(v_i, v_j) = 2$ if u_i is not adjacent to u_j and v_i is not adjacent to v_j . In order to satisfy this properties in this case, $|W_1| = \dim_{ls}(K_1 + G)$ while $|W_2| = \dim_{ls}(H)$ because $diam(H) \leq 2$. So, $\dim_{ls}(G + H) = \dim_{ls}(K_1 + G) + \dim_{ls}(H)$ if $diam(G) > 2$ and $diam(H) \leq 2$.

Case 3. If $diam(G) \leq 2$ and $diam(H) > 2$

According to Lemma 3.2, $d(u_i, u_j) = d(v_i, v_j) = 2$ if u_i is not adjacent to u_j and v_i is not adjacent to v_j . In order to satisfy this properties in this case, $|W_1| = \dim_{ls}(G)$ because $diam(G) \leq 2$ while $|W_2| = \dim_{ls}(K_1 + H)$. So, $\dim_{ls}(G + H) = \dim_{ls}(G) + \dim_{ls}(K_1 + H)$ if $diam(G) \leq 2$ and $diam(H) > 2$.

Case 4. If $diam(G) \leq 2$ and $diam(H) \leq 2$ or $\dim_{ls}(G) = \dim_{ls}(H) = 1$

Since $diam(G) \leq 2$ and $diam(H) \leq 2$ then $|W_1| = \dim_{ls}(G)$ and $|W_2| = \dim_{ls}(H)$. Assume $\dim_{ls}(G + H) = \dim_{ls}(G) + \dim_{ls}(H)$. Let $u_k \in W_1$ and $v_k \in W_2$, then there is a pair of adjacent vertex (u_i, v_j) where $u_i \notin W_1, v_j \notin W_2, u_i$ adjacent with u_k , and v_j adjacent with v_k that applies $u_i \notin I[v_j, u_k], u_i \notin I[v_j, v_k], v_j \notin I[u_i, u_k],$ and $v_j \notin I[u_i, v_k]$. So, $\dim_{ls}(G + H) \neq \dim_{ls}(G) + \dim_{ls}(H)$. Following Lemma 3.1, $\dim_{ls}(G + H) = \dim_{ls}(G) + \dim_{ls}(H) + 1$ if $diam(G) \leq 2$ and $diam(H) \leq 2$ or $\dim_{ls}(G) = \dim_{ls}(H) = 1$. ■

Using theorems in Section 1, 2, and 3 we can get the local strong metric dimension of $G + H$ if G and H are two of the following graphs, which are path (P_n), cycle (C_n), complete graph (K_n), and star graph (S_n). Since $diam(P_n) \geq 2$, $diam(C_n) \geq 2$ for $n > 5$ and $diam(K_n) = 1 \leq 2$, $diam(S_n) = 2$ then according to Theorem 3.1:

- i. $\dim_{ls}(P_n + P_m) = \dim_{ls}(K_1 + P_n) + \dim_{ls}(K_1 + P_m)$ if $n, m > 5$
- ii. $\dim_{ls}(P_n + C_m) = \dim_{ls}(K_1 + P_n) + \dim_{ls}(K_1 + C_m)$ if $n, m > 5$
- iii. $\dim_{ls}(P_n + K_m) = \dim_{ls}(K_1 + P_n) + \dim_{ls}(K_m)$
- iv. $\dim_{ls}(P_n + S_m) = \dim_{ls}(K_1 + P_n) + \dim_{ls}(S_m)$
- v. $\dim_{ls}(C_n + C_m) = \dim_{ls}(K_1 + C_n) + \dim_{ls}(K_1 + C_m)$ if $n, m > 5$
- vi. $\dim_{ls}(C_n + K_m) = \dim_{ls}(K_1 + C_n) + \dim_{ls}(K_m)$
- vii. $\dim_{ls}(C_n + S_m) = \dim_{ls}(K_1 + C_n) + \dim_{ls}(S_m)$
- viii. $\dim_{ls}(K_n + K_m) = \dim_{ls}(K_n) + \dim_{ls}(K_m) + 1$
- ix. $\dim_{ls}(K_n + S_m) = \dim_{ls}(K_n) + \dim_{ls}(S_m) + 1$
- x. $\dim_{ls}(S_n + S_m) = \dim_{ls}(S_n) + \dim_{ls}(S_m) + 1$

4. Conclusion

In this paper we get the result regarding local strong metric dimension in the join of graphs $G + H$ which is:

$$\dim_{ls}(G + H) = \begin{cases} \dim_{ls}(K_1 + G) + \dim_{ls}(K_1 + H) & \text{if } diam(G) > 2 \text{ and } diam(H) > 2 \\ \dim_{ls}(K_1 + G) + \dim_{ls}(H) & \text{if } diam(G) > 2 \text{ and } diam(H) \leq 2 \\ \dim_{ls}(G) + \dim_{ls}(K_1 + H) & \text{if } diam(G) \leq 2 \text{ and } diam(H) > 2 \\ \dim_{ls}(G) + \dim_{ls}(H) + 1 & \text{if } diam(G) \leq 2 \text{ and } diam(H) \leq 2 \\ & \text{or } \dim_{ls}(G) = \dim_{ls}(H) = 1 \end{cases}$$

In the next research, we suggest other researchers to develop the concept of local strong metric dimension into another concept of metric dimension, for example local strong complement metric dimension.

Acknowledgement

Thank you to Mathematics Department, Mathematics and Science Faculty, and LPPM Universitas Islam Madura that has provided support in this research.

References

- [1] Adawiyah R, Dafik, Alfarisi R, Prihandini R M, Agustin I H and Venkatachalam M 2020 *J. of Phys: Conf. Series* **1543** 1-7
- [2] Amri Z, Irvan I, Maryanti I and Sumardi H 2021 *J. of Phys: Conf. Ser.* **1731** 1-5
- [3] Cahyabudi A N and Kusmayadi T A 2017 *IOP Conf. Ser.: J. of Phys* **909** 1-5
- [4] Ganguli R and Roy S 2017 *Int. J. of Comp. and Applied Math.* **12** 469-85
- [5] Kuziak D, Yero I G, and Velázquez J A R 2013 *Discrete Applied Math.* **161** 1022-27
- [6] Mintarsih Y and Kusmayadi T A 2018 *IOP Conf. J. of Phys: Conf. Ser.* **1008** 1-6
- [7] Okamoto F, Crosse L, Phinezy B, Zhang P and Kalamazoo 2010 *Mathematica Bohemica* **135** 239-55
- [8] Raza H, Liu J B, and Qu S 2020 *Access* **8** 11560-69
- [9] Susilowati L, Rosfiana A and Purwati U D 2018 *Discrete Applied Math.* Manuscript Number DA9697, submitted.
- [10] Sutardji N A, Susilowati L and Purwati U D 2019 *Contemporary Math. and App.* **1** 64-73

The local strong metric dimension in the join of graphs

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Ruzika Rimadhany, Darmaji. "Local metric dimension of circulant graph $\text{circ}(n:1,2,\dots,n+12)$ ", AIP Publishing, 2017
Publication 5%
- 2** Submitted to Universitas Nahdlatul Ulama Surabaya
Student Paper 5%
- 3** YINGYING ZHANG, XIAOYU ZHU. "Proper Vertex Connection and Graph Operations", Journal of Interconnection Networks, 2019
Publication 4%

Exclude quotes On

Exclude matches < 3%

Exclude bibliography On